

REGIONAL ANAESTHESIA

Hypnosis and communication reduce pain and anxiety in peripheral intravenous cannulation: Effect of Language and Confusion on Pain During Peripheral Intravenous Catheterization (KTHYPE), a multicentre randomised trial

Nicolas Fusco^{1,2}, Franck Bernard^{2,3}, Fabienne Roelants^{2,4}, Christine Watremez^{2,4}, Hervé Musellec^{2,3}, Bruno Laviolle⁵, Helene Beloeil^{1,*} on behalf of the Effect of Language and Confusion on Pain During Peripheral Intravenous Catheterization (KTHYPE) group^{†,‡}

¹Anesthesia and Intensive Care Department, Centre Hospitalier Universitaire de Rennes, Université de Rennes, Inserm CIC 1414 Numecan, Rennes, France, ²Émergences Campus, Hypnosis Institute, Rennes, France, ³Anesthesia Department, Centre Hospitalier privé Saint Grégoire, Saint Grégoire, France, ⁴Anesthesiology Department, Cliniques universitaires Saint-Luc, Université catholique de Louvain, Brussels, Belgium and ⁵Clinical Pharmacology Department, Centre Hospitalier Universitaire de Rennes, Université de Rennes, Inserm, Rennes, France

*Corresponding author. E-mail: helene.beloeil@chu-rennes.fr

[†]Mustapha Ahmin, Sébastien Barthes, Natacha Boulon, Patricia Branchu, Catherine Copin-Eliat, Séverine Delahaye-Larralde, Pierre Fillatre, Charlotte Donal, Pierre Hugot, Cécile Gauvrit, Nicolas Guillou, Isabelle Leroyer, Cassiopée Pouliquen, Alain Renault, Emilie Rolland, Chloé Rousseau, Julie Rullier, Simon Schonne, and Thomas Yven.

[‡]The results of this study were presented at the annual meeting of the European Society of Anaesthesiology in Copenhagen (June 2018) and at the annual meeting of the French Society of Anesthesia & Intensive Care Medicine in Paris (September 2018).

Abstract

Background: Clinicians traditionally warn patients of pain before peripheral i.v. cannulation (PIVC). However, using words related to pain or undesirable experiences can result in greater pain and anxiety. The use of positive words can improve pain perception and subjective patient experience. We aimed to compare the effects of three types of communication, including hypnotic communication, on pain, comfort, and anxiety in patients during PIVC.

Methods: The Effect of Language and Confusion on Pain During Peripheral Intravenous Catheterization (KTHYPE) trial is a randomised, parallel, single-blind, multicentre study of patients undergoing PIVC on the dorsal face of the hand before surgery. Patients from three hospitals were randomly allocated to one of three groups: PIVC performed with a hypnosis technique (hypnosis group), negative connotation (nocebo group), and neutral connotation (neutral group). The primary outcome measure was the occurrence of pain measured with a 0–10 numerical rating scale just after PIVC.

Results: Of the 272 subjects analysed (hypnosis, $n=89$; nocebo, $n=92$; neutral, $n=91$), pain after PIVC was lower in the hypnosis group (mean [standard deviation]; range) (1.5 [1.9]; 0–5) compared with the neutral (3.5 [2.3]; 0–9; $P<0.0001$) and nocebo groups (3.8 [2.5]; 0–10; $P<0.0001$). Whilst anxiety was higher and comfort lower before PIVC in the hypnosis group, anxiety decreased and comfort perception increased after PIVC when hypnosis was used.

Conclusions: This is one of the first well-designed RCTs showing a significant benefit of a hypnosis technique during a routine procedure, such as PIVC. The results could facilitate implementation of hypnosis in daily clinical care.

Clinical trial registration: NCT02662322.

Keywords: anaesthesia; anxiety; hypnosis; pain; patient experience; peripheral intravenous cannulation

Editor's key points

- The use of positive words can improve pain perception and subjective patient experience.
- The Effect of Language and Confusion on Pain During Peripheral Intravenous Catheterization (KTHYPE) trial was a randomised, parallel, single-blind, multicentre study comparing the effects of three types of communication on pain, comfort, and anxiety in preoperative patients during peripheral i.v. cannulation (PIVC).
- Pain and anxiety were decreased and comfort perception increased after PIVC when positive communication and hypnosis were used.
- This RCT showed a significant benefit of a hypnosis technique during a routine perioperative procedure.

Communication is decisive in creating a therapeutic alliance with the patient. Unfortunately, a routine procedure as frequent and simple as peripheral i.v. cannulation (PIVC) can be a source of stress and pain. Pharmacological interventions have been shown to reduce the pain associated with PIVC. A network meta-analysis suggested that PIVC pain can be reduced by local anaesthesia.¹ However, local anaesthesia for cannulation is usually only offered to children, and fewer than half of clinicians follow this procedure for adults,^{2,3} which is time-consuming³ and has a high rate of puncture failure.^{2,4} Psychological interventions have also been shown to reduce the pain and anxiety associated with PIVC in children.^{5–7}

Communication skills are recognised as a key element of care,⁸ but unfortunately, clinicians often warn adult patients of pain using words with a negative connotation (i.e. 'painful' and 'sting'). This attitude is thought to be helpful and empathic. However, warning using language that refers to negative experiences is associated with a modification of pain and comfort perception.^{9,10} Warning the patient that the act will be painful leads to more pain and anxiety, and can create a nocebo effect.^{9–13} On the contrary, the use of positive words and sentences can benefit patient comfort.^{14,15} Hypnotic communication uses positive suggestions and distractions to reduce the sensitive and affective dimensions of the pain experience. Hypnosis is a state of consciousness involving focused attention and reduced peripheral awareness characterised by an enhanced capacity for response to suggestion.¹⁶

Indirect hypnosis or Ericksonian hypnosis is a method utilising body language, conversation, metaphors, and other hypnosis techniques to induce a therapeutic behavioural change by indirect suggestion. It starts from the beginning of the relationship between the therapist and the patient. This therapeutic alliance is essential. The hypnotic confusion technique uses distraction to focus patients on an incongruous, unexpected element outside of their preoccupation of the moment. By distracting the conscious mind, the therapist is able to open the unconscious mind to hypnotic language and to take advantage of its induced suggestibility to deliver an indirect suggestion of comfort. A benefit of hypnosis has been shown for anxiety and pain,^{17–20} but only few methodologically rigorous studies applying minimally effective control conditions have been published.^{17,21} We therefore assessed the effects of hypnosis using positive words associated with a

confusion technique on the subjective experience of PIVC when compared with a nocebo and neutral communication.

Methods**Study design and population**

The Effect of Language and Confusion on Pain During Peripheral Intravenous Catheterization (KTHYPE) study is a randomised, controlled, parallel-group, single-blind, multicentre, international clinical trial conducted in three hospitals: Rennes University Hospital (Rennes, France); Saint-Grégoire Private Hospital (Saint Grégoire, France); and Cliniques universitaires Saint-Luc, Université catholique de Louvain (Brussels, Belgium). The Rennes University Hospital Institutional Review Board reviewed and approved this clinical investigation (No. ID-RCB 2015-A01353-46), which was registered at <https://clinicaltrials.gov/ct2/show/NCT02662322> (NCT02662322). The study was approved by all local ethics committees (No. 2015-A01353-46–2016 01 07).

Adult patients >18 yr of age requiring a 20G PIVC on the dorsal surface of the hand before a scheduled surgery were included. The non-inclusion criteria were unable to communicate in French, history of difficult venous access, premedication, pregnant or breastfeeding women, legally protected (under judicial protection, guardianship or supervision, and persons deprived of their liberty), and urgent surgery. In case of failure during the first attempt of PIVC, the patient would be excluded from the study. All eligible patients gave their written consent before study participation.

Procedure

An information sheet about the study was given to all patients during the preoperative consultation. After arriving in the operating room, eligible patients were informed of the study by an anaesthesiologist not involved in the care of the patient and is called an 'anaesthesiologist researcher'. He/she was blind to the allocation group and proposed to the patient to participate as follows: 'A peripheral intravenous cannulation is necessary for your anaesthesia. We currently investigate how different procedures are perceived during the venous cannulation process. If you accept to participate in this trial, we will ask you to evaluate your experience during venous cannulation'.

The subjects were randomised into three groups (hypnosis, nocebo, and neutral) using a computer-generated randomisation table stratified by an investigation site with a 1:1:1 ratio. The subjects were blind of the allocation group. The primary evaluation criterion was assessed by the patient him-/herself (self-evaluation). The anaesthesiologist researcher assessed the presence of movements or vocalisation during PIVC. During the study period, the subjects and anaesthesiologist researchers were kept blind to the randomisation group. The anaesthesiologist and the anaesthesiologist nurse, both called clinicians in the study, were different from the anaesthesiologist researcher and were not blinded. They did not participate in the assessment of the subjects at any time.

In all groups, the clinician's experience for PIVC was at least 3 yr. No local anaesthesia was allowed before PIVC. Only one attempt was allowed, and the clinician catheterising was the only one talking to the patient at the time. In the hypnosis group, clinicians were anaesthesiologists or nurses with a diploma in therapeutic and hypnotic communication, and a

Table 1 Structured and standardised communication during peripheral i.v. cannulation.

	Placing a tourniquet around the arm	Decontaminating the skin with light friction	Inserting a peripheral i.v. catheter
Hypnosis	'How did you come to the hospital?'	'How long did it take you to come here?'	'Is your bike still going to the pool?'
Neutral	'I am placing the tourniquet to dilate the vein'.	'I am decontaminating the skin'.	'I am putting the catheter in'.
Nocebo	'I am placing the tourniquet; it grips/squeezes hard!'	'It is cold!'	'Warning, I will sting. One, two, three, it stings!'

minimum of 1 yr of experience. In the nocebo and neutral groups, clinicians were anaesthesiologists or nurses with no training in therapeutic communication and hypnosis. During the procedure, the clinicians communicated with the patient in a structured and standardised way depending on the allocation group (Table 1). In the hypnosis group, the clinicians applied classical non-verbal hypnotic tools adapted to the subject and indirect suggestion of comfort by body language. The whole procedure lasted <5 min.

Outcomes

The primary outcome was occurrence of pain just after PIVC, assessed using a numerical rating scale (NRS).²² The secondary outcomes were perception of comfort and anxiety before and after PIVC measured with a validated NRS.^{23,24} Pain, comfort, and anxiety were self-evaluated by the subjects minutes before and after PIVC. The subjects evaluated themselves before seeing any needle.

Data collection

At each participating centre, data were collected and entered into the electronic web-based case report form by two investigators (NF and FR) blinded to the allocation group, under the supervision of the trial site investigators. Data collection was monitored by trained clinical research associates.

Data collected by the anaesthesiologist researcher were anthropometric (age, gender, and BMI), type of surgical procedures according to the classification of the American College of Cardiology/American Heart Association,²⁵ standard of education, and socio-economic category. To ensure a blinded evaluation, the presence or not of spontaneous patient arm, face withdrawal, smile, or an unprompted vocalisation or comments were also recorded by the anaesthesiologist researcher.

The subjects were asked to quantify their pain, anxiety, and comfort on 11-point NRS (0=no pain to 10=worst imaginable pain experience, 0=no anxiety to 10=worst imaginable anxiety, and 0=no comfort to 10=best imaginable comfort) just before (for anxiety and comfort only) and within 3 min after the completed PIVC.

Statistical analysis

Based on an expected mean (standard deviation [SD]) NRS value of 4.4 (2.4) in the nocebo,¹³ we included 88 subjects in each group to show a decrease in NRS of 15% in the neutral group and 30% in the hypnosis group (effect size of 0.05), assuming an alpha risk of 5% and a power of 90% in a two-sided one-way analysis of variance (nQuery 8, version 8.3.1.0; Statsols, Cork,

Ireland). Based on Dutt-Gupta and colleagues,¹³ who calculated the sample size with a hypothesised 20% decrease in NRS, we considered that a 30% decrease in NRS would reflect a clinically relevant effect, and that 15% would indicate a significant but less clinically relevant effect in the neutral group compared with nocebo. We included 10% more subjects to take into account expected cases of failure of the first attempt of PIVC, or 300 subjects to obtain a minimum of 264 included.

Statistical analysis was performed with SAS version 9.4 (SAS Institute, Cary, NC, USA). All data analyses were performed by researchers blinded to the group allocation. Normal distribution was assessed by descriptive statistics and histograms. Quantitative variables are presented as mean (SD) for continuous variables and as *n* (%) for categorical variables. An analysis of study outcomes was performed by one-way analysis of variance for continuous variables, and χ^2 test or Fisher's exact test for categorical variables. In cases of statistical significance, two-by-two comparisons were performed with a Bonferroni adjustment threshold for multiple testing. A sensitivity analysis taking into account the standard of education was also performed for the primary outcome using two-way analysis of variance, in which the group–standard of education interaction was tested.

Results

Between March 2016 and March 2017, a total of 294 patients were randomised (Fig. 1) from Rennes University Hospital: 62 recruited/50 analysed; Saint-Grégoire Private Hospital: 132 recruited/128 analysed; and Cliniques universitaires Saint-Luc, Université catholique de Louvain: 100 recruited/94 analysed. The first attempt cannulation failure was 6.5%, leaving 272 subjects analysed (hypnosis group, *n*=89; neutral group, *n*=92; nocebo group, *n*=91). The subject characteristics are presented in Table 2. The study groups were similar in gender, BMI, types of surgical procedures, standard of education, or socio-economic category. Self-evaluation of pain, anxiety, and comfort was performed within 3 min after the end of PIVC (2.5 [2.8] min) without differences between groups.

The primary outcome, pain after PIVC, was significantly lower in the hypnosis group compared with the neutral and nocebo groups (Fig. 2). This difference in pain after PIVC was still significant after adjustment for standard of education. There was no significant difference in pain after PIVC between the neutral and nocebo groups.

The subjects in the hypnosis group were less likely to spontaneously withdraw their arm or face (*n*=1 [2%]) compared with the neutral group (*n*=17 [31.5%]) or nocebo group (*n*=15 [22.4%]) (*P*<0.05 for both comparisons without difference between neutral and nocebo groups [*P*=0.78]). Negative face (grimace in pain) was higher in the nocebo group




Fig 1. Effect of Language and Confusion on Pain During Peripheral Intravenous Catheterization (KTHYPE) study flow chart.

($n=32$ [47.8%]; $P<0.01$) and neutral group ($n=27$ [50%]; $P<0.01$) compared with the hypnosis group ($n=9$ [18%]). Positive face (smile) was higher in the hypnosis group ($n=31$ [62%]) compared with the nocebo ($n=2$ [3%]; $P<0.0001$) and neutral ($n=1$ [2%]; $P<0.0001$) groups. No difference between groups was observed for unprompted vocalisation or comments.

Anxiety before PIVC was higher in the hypnosis group compared with the neutral group, but not with the nocebo group (Table 3). Comfort before PIVC was lower in the hypnosis group compared with the neutral and nocebo groups. Anxiety after PIVC decreased in the hypnosis group, but not in the nocebo and neutral groups. Comfort increased after PIVC in the hypnosis group, and decreased in the neutral and nocebo

groups. When anxiety and comfort after PIVC were adjusted to those before PIVC, anxiety after PIVC was significantly lower and comfort was significantly higher in the hypnosis group compared with the neutral and nocebo groups (Fig. 3). No differences were observed between the neutral and nocebo groups.

Discussion

To our knowledge, this is the first RCT showing the benefit of hypnosis on a routine procedure as simple as PIVC. Pain and anxiety decreased and comfort perception increased after PIVC when hypnosis was used. Previous studies reported pain

Table 2 Subject characteristics. There were no significant differences ($P<0.05$) between study groups. ACC/AHA, American College of Cardiology/American Heart Association; SD, standard deviation.

	Total (n=272)	Hypnosis (n=89)	Neutral (n=92)	Nocebo (n=91)
Demographics				
Mean age (range) (yr)	54.5 (18 – 89)	56.2 (22 – 89)	53.7 (18 – 86)	53.7 (19 – 87)
Sex (F/M) (%)	55/45	54/46	52/48	60/40
Mean BMI (SD) (kg m^{-2})	25.7 (5.2)	26 (5.9)	25.1 (4.3)	25.9 (5.3)
Surgery procedure risk (ACC/AHA), n (%)				
High	11 (4)	8 (9)	2 (2)	1 (1)
Intermediate	81 (30)	22 (25)	27 (29)	32 (35)
Low	180 (66)	59 (66)	63 (68)	58 (64)
Standard of education, n (%)				
No qualification	26 (10)	10 (11.4)	11 (12.0)	5 (5.5)
School (USA)/secondary school (UK)	75 (28)	24 (27.3)	29 (31.5)	22 (24.2)
High school (USA)/A levels (UK)	60 (22)	16 (18.2)	19 (20.7)	25 (27.5)
University (USA)/higher education (UK)	110 (41)	38 (43.2)	33 (35.9)	39 (42.9)




Fig 2. Self-evaluated pain after peripheral i.v. cannulation on an 11-point numerical rating scale (from 0=no pain to 10=worst imaginable pain experience) for the hypnosis ($n=89$), neutral ($n=92$), and nocebo ($n=91$) groups. Median, first quartile, and third quartile are represented by boxes. Whisker plots represent 1.5 inter-quartile space associated with outliers for the hypnosis group. Mean is represented by rhombs. *** $P<0.0001$. NS, not significant.

scores after 20G PIVC was placed on the dorsal surface of the hand similar to values observed in our control groups (neutral/nocebo).^{26–28} The levels of pain NRS obtained with hypnotic communication in our study were similar to those reported with local anaesthesia.^{26–28} Therefore, hypnosis with confusion technique seems to offer a benefit comparable with invasive pharmacological interventions. Moreover, this benefit was not associated with side-effects and was not time-consuming (<3 min). Our first attempt failure rate (6.5%) was lower than in previous studies (7–16%).^{29,30} The low rate of failure in our study could be explained by the lack of local

anaesthesia, which can increase puncture failure,^{28,30} and by the experience of the staff in our study. Our results also show that a hypnotic confusion technique could be applied without impacting the efficiency of PIVC. A previous study showed the influence of negative words on pain during blood sampling.³¹ However, the physicians involved in that study were not trained in hypnotic communication. The non-verbal part of the experience was then missing. In our study, only the verbal language could be standardised. Our therapists adapted complex non-verbal communication to each subject, and indirect suggestions were induced from the outset with the therapeutic alliance.

Dutt-Gupta and colleagues¹³ showed that warning patients of a ‘sting’ before PIVC may not be helpful. Comparing communication with positive or negative words, they reported no differences in pain and Likert scale scores. However, they reported (as a secondary outcome) less patients vocalising pain during PIVC with positive communication. In our study, hypnotic communication produced decreased pain perception not only compared with the nocebo group, but also with the neutral group. As the hypnotic process can play a crucial role in the modulation and perception of pain, our results show that the use of positive words involving hypnotic communication could also reduce anxiety and improve comfort. Furthermore, pain perception, anxiety, and comfort were similar in the neutral and nocebo groups, suggesting that a neutral attitude is as deleterious as a nocebo one. Our findings confirm that warning patients with nocebo and even neutral words, although made with good intent, induced discomfort. Hypnosis can be defined as an altered conscious state of focused attention that involves absorption, some dissociative elements, and an increased responsiveness to suggestion. In other words, the hypnotic cerebral process is well known to improve suggestibility. The verbal distraction and focalisation included in the hypnotic confusion technique by saying something like, ‘is your bike still going to the pool?’ just before PIVC created confusion in a patient’s mind. The patient may ask, ‘what did he/she say?’ and at this time the patient is ‘dissociated’ and focused on the meaning of the sentence rather than on the PIVC.

Overall patient satisfaction is correlated with communication and pain management,³² and can be improved with hypnosis.³³ In the perioperative setting, PIVC is one of the major sources of preoperative anxiety.³⁴ Comfort obtained by hypnotic communication before surgery may help improve patient satisfaction.¹⁷ Hypnotic communication, or at least therapeutic communication, should be mandatory in the initial training of caregivers.

One limitation is that the study was single blinded. We chose to perform PIVC in the hypnotic group by non-blinded anaesthesiologists and nurses with a hypnotic communication diploma. We hypothesised that hypnotic communication would have too much influence on non-verbal communication to not impact the relation with patients. If the clinicians with a hypnotic communication diploma are able to suggest comfort, they could also suggest pain and discomfort if they performed PIVC for the neutral and nocebo groups. Another limitation is the restriction of the hypnotic group to anaesthesia providers with a hypnotic communication diploma. Whether the salutary effects of hypnosis are the same when providers not trained in hypnosis use the same technique needs further research.

Our study did not show the benefit of the hypnotic confusion technique *per se*, as it was provided by clinicians with a

Table 3 Numerical rating scale (NRS) of anxiety and comfort self-evaluated before and after peripheral intravenous cannulation in the hypnosis group compared with the neutral and nocebo groups. NRS, mean (standard deviation) [range]. Comparisons of pain, anxiety, and comfort NRS after PIVC are adjusted to those before. * $P<0.05$ vs neutral. † $P<0.05$ vs nocebo. ‡ $P<0.05$ before vs after.

	Hypnosis ($n=89$)	Neutral ($n=92$)	Nocebo ($n=91$)
Before peripheral intravenous cannulation			
Anxiety	4.4 (2.6) [0–10]*	3.0 (2.6) [0–10]	3.5 (2.7) [0–10]
Comfort	7.5 (2.1) [2–10]*, †	8.5 (1.6) [3–10]	8.3 (1.8) [3–10]
After peripheral intravenous cannulation			
Anxiety	2.3 (2.5) [0–9]*, †	3.0 (2.9) [0–10]	3.6 (2.7) [0–10]
Comfort	8.5 (1.7) [2–10]*, †	7.7 (2.2) [1–10]	7.2 (2.1) [1–10]
Pain	1.5 (1.9) [0–9]*, †	3.5 (2.3) [0–9]	3.8 (2.5) [0–10]
Difference before and after peripheral intravenous cannulation			
Anxiety	−2.1 (2.9)‡	+0.0 (2.3)	+0.1 (2.3)
Comfort	+1.0 (2.1)‡	−0.9 (1.9)‡	−1.1 (2.1)‡




Fig 3. Self-evaluated anxiety and comfort after peripheral i.v. cannulation (PIVC) on an 11-point numerical rating scale for the hypnosis ($n=89$), neutral ($n=92$), and nocebo ($n=91$) groups. Median, first quartile, and third quartile are represented by boxes. Mean is represented by rhombs. Whisker plots represent 1.5 inter-quartile range associated with outliers. *** $P<0.0001$. NS, not significant.

hypnotic communication diploma. We cannot conclude that the hypnotic confusion technique provided by the clinicians without a diploma would have any benefit. Indeed, Lang and colleagues³⁵ showed that hypnosis, compared with attentive behaviour alone, provides greater effects on pain and anxiety reduction during invasive medical procedures. Therefore, hypnotic communication involving a confusion technique may be more efficient than distraction alone to reduce pain and anxiety during PIVC. Further studies are necessary to test this hypothesis. Finally, anxiety was more important in the hypnosis group before PIVC compared with the other groups. These differences happened despite randomisation. However, the subjects in the hypnosis group were less anxious and more comfortable after PIVC, which is what matters because it is what they will remember. When patients are in a comfortable state, they are likely to stay in that state and uncontrollable anxiety is prevented.³⁶ In other words, a good experience with PIVC can determine the quality of the subsequent experience in the hospital.

In conclusion, the KTHYPE trial is the first randomised multicentre study evaluating the effect of communication on pain, comfort, and anxiety in surgical patients undergoing PIVC. It shows that hypnotic communication with a confusion technique compared with neutral or nocebo communication decreases pain and anxiety after PIVC. These results suggest that implementation of hypnosis into daily care could lead to significant changes in the standard of care in anaesthesia.

Authors' contributions

Study conception/design: all authors

Critical input for methodology: BL

Statistical analysis: BL

Writing of first draft: NF

Writing of manuscript: HB

Critical revision/modification of manuscript: NF, FB, FR, CW, HM, HB

NF was the principal investigator. NF, FB, FR, CW, and HM included patients in their centres. All authors approved the final version of the manuscript.

Acknowledgements

The committee allowing grants in the teaching hospital of Rennes had no role in the design and conduct of the study; the collection, management, analysis, and interpretation of the data; and the preparation, review, and approval of the manuscript.

Declaration of interest

The authors declare that they have no conflicts of interest.

Funding

Teaching hospital of Rennes, France

References

- Bond M, Crathorne L, Peters J, et al. First do no harm: pain relief for the peripheral venous cannulation of adults, a systematic review and network meta-analysis. *BMC Anesthesiol* 2016; **16**: 81
- Sado DM, Deakin CD. Local anaesthesia for venous cannulation and arterial blood gas sampling: are doctors using it? *J R Soc Med* 2005; **98**: 158–60
- Norris WD. The use of local anaesthesia in peripheral venous cannulation: current practice of junior doctors. *J R Nav Med Serv* 2002; **88**: 62–4
- Hendry F, Checketts MR, McLeod GA. Effect of intradermal anaesthesia on success rate and pain of intravenous cannulation: a randomized non-blind crossover study. *Scott Med J* 2011; **56**: 210–3
- Birnie KA, Noel M, Chambers CT, Uman LS, Parker JA. Psychological interventions for needle-related procedural pain and distress in children and adolescents. *Cochrane Database Syst Rev* 2018; **33**: 164
- Koller D, Goldman RD. Distraction techniques for children undergoing surgery: a critical review of pediatric research. *J Pediatr Nurs* 2012; **27**: 652–81
- Liossi C, White P, Hatira P. A randomized clinical trial of a brief hypnosis intervention to control venepuncture-related pain of paediatric cancer patients. *Pain* 2009; **142**: 255–63
- Smith AF, Mishra K. Interaction between anaesthetists, their patients, and the anaesthesia team. *Br J Anaesth* 2010; **105**: 60–8
- Lang EV, Hatzopoulos O, Koch T, et al. Can words hurt? Patient-provider interactions during invasive procedures. *Pain* 2005; **114**: 303–9
- Bejenke CJ. Painful medical procedures. In: Barber J, editor. *Hypnosis and suggestion in the treatment of pain: a clinical guide*. London: W.W. Norton; 1996. p. 209–61
- Benedetti F, Carlino E, Pollo A. How placebos change the patient's brain. *Neuropsychopharmacology* 2011; **36**: 339–54
- Benedetti F, Lanotte M, Lopiano L, Colloca L. When words are painful: unraveling the mechanisms of the nocebo effect. *Neuroscience* 2007; **147**: 260–71
- Dutt-Gupta J, Bown T, Cyna AM. Effect of communication on pain during intravenous cannulation: a randomized controlled trial. *Br J Anaesth* 2007; **99**: 871–5
- Varellmann D, Pancaro C, Cappiello EC, Camann WR. Nocebo-induced hyperalgesia during local anesthetic injection. *Anesth Analg* 2010; **110**: 868–70
- Cyna AM, Andrew M. Words can help! *Pain* 2005; **117**: 239–40
- Elkins GR, Barabasz AF, Council JR, Spiegel D. Advancing research and practice: the revised APA Division 30 definition of hypnosis. *Int J Clin Exp Hypn* 2015; **63**: 1–9
- Kendrick C, Sliwinski J, Yu Y, et al. Hypnosis for acute procedural pain: a critical review. *Int J Clin Exp Hypn* 2016; **64**: 75–115
- Kekecs Z, Nagy T, Varga K. The effectiveness of suggestive techniques in reducing postoperative side effects: a meta-analysis of randomized controlled trials. *Anesth Analg* 2014; **119**: 1407–19
- Boly M, Faymonville M-E, Schnakers C, et al. Perception of pain in the minimally conscious state with PET activation: an observational study. *Lancet Neurol* 2008; **7**: 1013–20
- Faymonville ME, Laureys S, Degueldre C, et al. Neural mechanisms of antinociceptive effects of hypnosis. *Anesthesiology* 2000; **92**: 1257–67
- Cheseaux N, de Saint Lager AJ, Walder B. Hypnosis before diagnostic or therapeutic medical procedures: a systematic review. *Int J Clin Exp Hypn* 2014; **62**: 399–424
- Price DD, McGrath PA, Rafii A, Buckingham B. The validation of visual analogue scales as ratio scale measures for chronic and experimental pain. *Pain* 1983; **17**: 45–56
- Chooi CSL, White AM, Tan SGM, Dowling K, Cyna AM. Pain vs comfort scores after Caesarean section: a randomized trial. *Br J Anaesth* 2013; **110**: 780–7
- Crandall M, Lammers C, Senders C, Savedra M, Braun JV. Initial validation of a numeric zero to ten scale to measure children's state anxiety. *Anesth Analg* 2007; **105**: 1250–3
- Poldermans D, Bax JJ, Boersma E, et al. Guidelines for pre-operative cardiac risk assessment and perioperative cardiac management in non-cardiac surgery. *Eur J Anaesth* 2010; **27**: 92–137
- Burke SD, Vercler SJ, Bye RO, Desmond PC, Rees YW. Local anesthesia before IV catheterization. *Am J Nurs* 2011; **111**: 40–5
- Vaghadia H, al-Ahdal OA, Nevin K. EMLA patch for intravenous cannulation in adult surgical outpatients. *Can J Anaesth* 1997; **44**: 798–802
- Armstrong P, Young C, McKeown D. Ethyl chloride and venepuncture pain: a comparison with intradermal lidocaine. *Can J Anaesth* 1990; **37**: 656–8
- Page DE, Taylor DM. Vapocoolant spray vs subcutaneous lidocaine injection for reducing the pain of intravenous cannulation: a randomized, controlled, clinical trial. *Br J Anaesth* 2010; **105**: 519–25
- Biro P, Meier T, Cummins AS. Comparison of topical anaesthesia methods for venous cannulation in adults. *Eur J Pain* 1997; **1**: 37–42
- Ott J, Aust S, Nouri K, Promberger R. An everyday phrase may harm your patients. *Clin J Pain* 2012; **28**: 324–8
- Boulding W, Glickman SW, Manary MP, Schulman KA, Staelin R. Relationship between patient satisfaction with inpatient care and hospital readmission within 30 days. *Am J Manag Care* 2011; **17**: 41–8
- Amraoui J, Pouliquen C, Fraisse J, et al. Effects of a hypnosis session before general anesthesia on postoperative outcomes in patients who underwent minor breast cancer surgery. *JAMA Netw Open* 2018; **1**: e181164–4
- Tee FY, Siew C, Low L, Matizha P. Patient perceptions and experience of pain, anxiety and comfort during peripheral intravenous cannulation in medical wards: topical anaesthesia, effective communication, and empowerment. *Int J Nurs Sci* 2015; **5**: 41–6
- Lang EV, Benotsch EG, Fick LJ, et al. Adjunctive non-pharmacological analgesia for invasive medical procedures: a randomised trial. *Lancet* 2000; **355**: 1486–90
- Arnsten AF. Stress weakens prefrontal networks: molecular insults to higher cognition. *Nat Neurosci* 2015; **18**: 1376–85

Handling editor: Hugh C Hemmings Jr